

PHYS 528 Homework #5

Due: Feb.28, 2019

Note that Feb.19-22 is UBC reading week so there will be no class on Feb.19 or 21.

1. Scalar expansions.

Consider a general theory of n real scalars ϕ_i ($i = 1, \dots, n$) with Lagrangian

$$\mathcal{L} = \frac{1}{2} \sum_{i=1}^n (\partial\phi_i)^2 - V(\phi_i) .$$

If $\{\phi_i = \langle\phi_i\rangle\}$ is a minimum of the potential, show that the fields $h_i(x) = \phi_i(x) - \langle\phi_i\rangle$ have canonical normalization and vanish at the minimum of the potential. Next, expand the potential about the minimum in a power series in h_i and show that $\partial^2 V / \partial\phi_i \partial\phi_j |_{\langle\phi\rangle}$ is the mass matrix for the scalars h_i .

2. SSB and NGBs.

- In the spontaneously broken global $U(1)$ theory discussed in the notes, work out the kinetic term and potential in terms of the new polar field variables we defined.
- Suppose we have the same theory but with $-\mu^2 \rightarrow +\mu^2$ in the potential. Is there still spontaneous symmetry breaking? What are the particle masses of all the real scalar degrees of freedom? *Hint: we really want canonical kinetic terms!*
- For the same theory, expanded around the vacuum $\langle\phi\rangle = e^{i\beta} v$, work out the mass matrix $\partial^2 V / \partial\phi_i \partial\phi_j |_{\langle\phi\rangle}$ in terms of the original field variables ϕ and ϕ^* and show that it has a zero determinant. Also, find $F_i^a(\langle\phi\rangle)$ for an infinitesimal phase rotation and show that it is a zero eigenvalue of this mass matrix.
Hint: treat ϕ and ϕ^ as independent degrees of freedom.*
- For the global $SU(2) \times U(1)$ -symmetric theory discussed in the notes, work out the full Lagrangian in terms of the new field variables

$$\begin{pmatrix} \phi_+ \\ \phi_0 \end{pmatrix} = \begin{pmatrix} (\phi_{+r} + i\phi_{+i})/\sqrt{2} \\ v + (\phi_{0r} + i\phi_{0i})/\sqrt{2} \end{pmatrix}$$

This choice corresponds to an expansion about the vacuum with $\alpha^a = \beta = 0$. What are the mass eigenvalues?

3. A semi-realistic Higgs.

- Starting from the global $SU(2) \times U(1)$ -symmetric theory discussed in the notes, elevate this to a theory that is invariant under local $SU(2) \times U(1)$ transformations by adding an appropriate set of vector gauge fields and couplings. What is the corresponding Lagrangian?
Hint: each gauge factor has its own gauge field and its own gauge coupling.

b) Work out the commutation relations of the modified set of generators \tilde{t} and $\{t_{G/H}^B\}$ discussed in the notes.

c) Expand this theory around the vacuum after making a nice choice of gauge. Find the masses of all the physical scalars, and make sure their kinetic terms are canonical.

Hint: for the choice of gauge, start with the field expansion used in the global $SU(2) \times U(1)$ theory discussed in the notes and then simplify it enormously by choosing a gauge in analogy to what was done in the gauged $U(1)$ theory discussed there. Talk to me if you aren't sure how to proceed.

d) Expand the gauge-covariant kinetic term of the scalar field to find mass terms for the vector fields.

Hint: remember that $t^a = \sigma^a/2$ and compute the covariant derivative $D_\mu\phi$ as a two-component column vector. Then use the fact that the gauge fields are real to simplify its square. Also, recall that any 2×2 symmetric matrix can be diagonalized by an orthogonal matrix, and this matrix can be built from the orthonormalized eigenvectors.