

PHYS 526 Homework #0

Due: optional (but recommended)

1. Natural Units

- What is one second in GeV units?
- What is one meter in GeV units?
- The LHC is trying to create new particles with masses M on the order of a TeV. On dimensional grounds, we expect the production cross section for such particles to go like $\sigma \sim 1/M^2$. What does this correspond to in femptobarns ($1 \text{ fb} = 10^{-15} \text{ b}$)?
- The mass scale that corresponds to Newton's constant $G_N = 6.67 \times 10^{-11} \text{ m}^3/\text{kg s}^2$ is called the Planck mass. What is its value in GeV units?
- The age of the Universe is about 13.7 billion years. Express this in GeV units and compare it to the Planck mass you found above.

2. Matrices and Indices

- Show that for any two $n \times n$ matrices, $(MN)^t = N^t M^t$.
- Show that δ_{ij} ($i, j = 1, \dots, n$) is the $n \times n$ identity matrix.
- Prove the cyclicity of the trace: $\text{tr}(AB) = \text{tr}(BA)$.
- Show that ϵ_{ijk} is cyclic:

$$\epsilon_{ijk} = \epsilon_{jki} = \epsilon_{kij} . \quad (1)$$

Use this to show that the triple product ($n = 3$) is also cyclic in the sense $a \cdot (b \times c) = c \cdot (a \times b) = b \cdot (c \times a)$.

- For $M = \sigma^1$ (the Pauli matrix) and $v^t = (1, 1)$, evaluate $\sum_j M_{ij} v_j$ and $\sum_i M_{ii} v_i$.

3. Relativistic Indices

- Show that $\eta_{\mu\nu} \Lambda^\mu_\lambda \Lambda^\nu_\kappa = \eta_{\lambda\kappa}$ implies that Λ leaves invariant the dot product of any pair of 4-vectors.
- Prove $\eta_{\nu\lambda} \eta^{\mu\kappa} \Lambda^\lambda_\kappa := \Lambda_\nu^\mu = (\Lambda^{-1})^\mu_\nu$.
- Objects with more than one Lorentz index are called tensors. Like vectors, we raise and lower their indices with η (e.g. $T^{\mu\nu} = \eta^{\mu\lambda} T_\lambda^\nu = \eta^{\mu\lambda} \eta^{\nu\kappa} T_{\lambda\kappa}$). Under Lorentz transformations, each index gets a power of Λ (e.g. $T^{\mu\nu} \rightarrow \Lambda^\mu_\lambda \Lambda^\nu_\kappa T^{\lambda\kappa}$).
 - Show that if we treat it as a tensor, $\eta^{\mu\nu} \rightarrow \eta^{\mu\nu}$ under Lorentz transformations.
 - A pair of tensor indices are said to be antisymmetric if $A^{\mu\nu} = -A^{\nu\mu}$. Show that if A is antisymmetric, $A^{\mu\nu} v_\mu v_\nu = 0$ for any vector v_μ .
 - Show that $T^{\mu\nu} u_\mu v_\nu$ is Lorentz invariant for any tensor T and vectors u, v .

4. Suppose we have the decay $A \rightarrow B + C$ for particles with masses m_a , m_b , and m_c . Compute the momenta of the decay products in the rest frame of the decaying particle.

5. Evaluate the integral

$$I_2 = \int d^3 p_b \int d^3 p_c \frac{1}{E_b E_c} \delta^{(4)}(P - p_b - p_c) ,$$

where $P = (M, \vec{0})$, $p_b^0 = E_b = \sqrt{m_b^2 + \vec{p}_b^2}$, and $p_c^0 = E_c = \sqrt{m_c^2 + \vec{p}_c^2}$.