

PHYS 528 Homework #2

Due: Feb.4, 2021

1. Work out the differential cross section $d\sigma/d(\cos\theta)$ for the process $e^+e^- \rightarrow \mu^+\mu^-$, where θ is the CM-frame angle between the incident electron and the outgoing muon. You may work in the limit that $p^2 \gg m_e^2$ and ignore the electron mass ($m_e \rightarrow 0$), but do keep the full dependence on the muon mass (instead of dropping it like we did in class). *Hint: note that the differential cross is what you get from the cross section formula but without doing the final integral over the outgoing angle θ .*
2. Compute the summed and squared matrix element for $e^-\mu^- \rightarrow e^-\mu^-$ scattering to leading order in QED at very high energy, $E_{CM} \gg m_\mu, m_e$. This implies that you can neglect the fermion masses.
(Optional: use this to compute the differential and total cross sections.)
3. Consider a massive Z' vector boson that couples to electrons with a vertex factor equal to $-ig'\gamma^\mu$.

- a) A massive vector has three independent polarization states. These can be represented by any three independent unit 4-vectors $\epsilon_\mu(p, \lambda)$ satisfying the constraints $p^\mu \epsilon_\mu = 0$ and $\epsilon_\mu^*(p, \lambda) \epsilon^\mu(p, \lambda') = -\delta_{\lambda\lambda'}$, where p^μ is the four-momentum of the vector boson and $\lambda = 1, 2, 3$ labels the three different polarizations. Find a simple set of polarization vectors in the rest frame of the massive vector. Show that they satisfy the constrained completeness relation

$$\sum_{\lambda} \epsilon_\mu(p, \lambda) \epsilon_\nu^*(p, \lambda) = -\eta_{\mu\nu} + p_\mu p_\nu / m_{Z'}^2.$$

- b) Compute the total unpolarized decay width for $Z' \rightarrow e^+e^-$. For this, use the vertex and the completeness relation stated above. Keep the full dependence on the masses of the Z' and the electron.